

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 July 2011 (14.07.2011)

(10) International Publication Number
WO 2011/084585 A2

(51) International Patent Classification:
C07D 221/12 (2006.01)

(21) International Application Number:
PCT/US2010/060735

(22) International Filing Date:
16 December 2010 (16.12.2010)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/287,115 16 December 2009 (16.12.2009) US

(71) Applicant (for all designated States except US): **WASHINGTON UNIVERSITY** [US/US]; One Brookings Drive, St. Louis, Missouri 63130 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MACH, Robert** [US/US]; 519 Meramec View Drive, Eureka, Missouri 63025 (US). **MINTUN, Mark** [US/US]; 2 Glazeview


Road, Town and Country, Missouri 63017 (US). **CHU, Wenhua** [US/US]; 10333 Garibaldi Place, Saint Louis, Missouri 63131 (US). **DUGAN, Laura** [US/US]; 11109 Corte Mar de Cristal, San Diego, California 92130 (US).

(74) Agent: **ZACKSON, Saul, L.**; Zackson Law LLC, 1015 Locust Street, Suite 750, Saint Louis, Missouri 63101 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

[Continued on next page]

(54) Title: DIHYDROETHIDINE ANALOGUES AND USES THEREOF



(57) Abstract: Tracers for imaging distribution of reactive oxygen species (ROS) are disclosed. The tracers include radiolabeled dihydroethidine (DHE) analogues. Further disclosed are uses of the compounds, including methods of imaging tissue distribution of ROS in vivo by positron emission tomography (PET). Methods of synthesizing the compounds are also disclosed.

WO 2011/084585 A2

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,

SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report (Rule 48.2(g))*

DIHYDROETHIDINE ANALOGUES AND USES THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority to U.S. Provisional Application Serial No. 61/287,115 filed on December 16, 2009, which is incorporated herein by reference in its entirety.

5

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

The disclosed subject matter was developed in part with Government support under grant R01AG036045 from the National Institutes of Health. The Government has certain rights in the invention.

10

INTRODUCTION

The present teachings relate to the field of free radicals in biology and medicine.

Free radicals play key roles in the pathogenesis of a large number of diseases and processes, for example brain processes related to Alzheimer's disease, other neurodegenerative diseases, severe mental illnesses, accelerated aging, and vascular disease. Free radicals are molecules containing one or more unpaired orbital electrons, which can increase the molecule's reactivity. Radicals of oxygen are a predominant class of free radicals in the human body. Free oxygen radicals in biology are termed reactive oxygen species (ROS). ROS include the superoxide anion radical, which is a predominant ROS in the brain. ROS are considered important in both acute and chronic inflammation pathways, and can affect multiple metabolic processes and signaling molecules in the brain. Imaging the distribution of ROS is of considerable interest to physicians and other health care professionals.

15

20

25


Dihydroethidine (DHE) is a molecule that can be oxidized by superoxide. When administered peripherally, DHE can enter the brain via the bloodstream and can be oxidized by superoxide. Once oxidized, DHE can be trapped behind the blood brain barrier, and can accumulate there. DHE has been used for visualizing ROS distribution in animal tissue using fluorescence techniques (Quick and Dugan, *Ann. Neurology*, 49:627-635, 2001; Garbett, N.C., et al., *Biophys. J.* 87: 3974-3981, 2004). However, health care professionals are in need of

tracers that can be used for positron emission tomography (PET) imaging of ROS distribution in vivo.


SUMMARY

The present inventors have developed a series of compounds which can be used as radiolabels for diagnostic imaging, in particular positron emission tomography (PET) imaging of ROS distribution in the tissue of a mammal such as a human. The compounds include analogues of dihydroethidium (DHE) (also known as 2,7-diamino-10-ethyl-9-phenyl-9,10-dihydrophenanthridine, 8-Diamino-5,6-dihydro-5-ethyl-6-phenylphenanthridine and hydroethidine). Analogues of DHE include compounds that are structurally similar to DHE but differ slightly in composition, such as in the replacement of one atom by an atom of a different element or in the presence of a particular functional group.

In some embodiments, the present teachings disclose a radiolabeled compound or salt thereof of structure




15 wherein R₁ can be H or O-R₃, R₃ can be (CH₂)_qCH₃, or (CH₂)_p-F, q can be an integer from 0 to 10; n can be an integer from 0 to 3, m can be an integer from 0 to 3,




and p can be an integer from 0 to 3, R_2 can be methyl or an integer from 0 to 10. In some configurations, q can be 0.


In further embodiments of the present teachings, the inventors disclose a radiolabeled compound or salt thereof of structure



5 wherein R_1 can be H or O- R_3 ; R_3 can be




10 $(CH_2)_q-CH_3$, and wherein q can be an integer from 0 to 10; n can be an integer from 0 to 3; m can be an integer from 0 to 3; p can be an integer from 0 to 3; and a can be an integer from 2 to 10. In some configurations, R_1 can be H. In some configurations, a CH_3 can be an $^{11}CH_3$. In some configurations, R_3 can be $(CH_2)_q-CH_3$ and q can be 0. In some configurations, R_3 can be $^{11}CH_3$. In some configurations, R_3 can be



, n can be an integer from 0 to 3, and m can be an integer from 0 to 3. In some

configurations, R_1 can be $O-R_3$, and R_3 can be $^{11}CH_3$. In some configurations, F can be ^{18}F .

In additional embodiments of the present teachings, the inventors disclose methods of imaging ROS in a mammal such as a human. These methods comprise administering to the mammal a radiolabeled compound or salt thereof of structure




wherein R_1 can be H or $O-R_3$, R_3 can be $(CH_2)_qCH_3$, , or , n can be an integer from 0 to 3, m can be an integer from 0 to 3, p can be integer from 0 to 3, and q can be an integer from 0 to 10, and subjecting the mammal to PET scanning.

In some embodiments, a compound of the present teachings can comprise a radioisotope such as a positron-emitting radioisotope, such as ^{18}F or ^{11}C . In some embodiments, a compound of the present teachings can be a radiolabeled compound or salt thereof as disclosed herein, wherein if R_1 is H, then the CH_3 can be $^{11}CH_3$.


In some embodiments, a compound of the present teachings can be a radiolabeled compound or salt thereof as disclosed herein, wherein if R_1 is $O-R_3$, and $q=0$, then R_3 can be $^{11}CH_3$.

In some embodiments, a compound of the present teachings can be a radiolabeled compound or salt thereof as disclosed herein, wherein if R_1 can be $O-R_3$, and R_3 can be a




fluorinated moiety such as wherein n can be an integer from 0 to 3, m can be an integer from 0 to 3, the F can be an ¹⁸F. In some configurations, n can be 1 and m can be 2.

In some embodiments, a compound of the present teachings can be a radiolabeled compound or salt thereof as disclosed herein, wherein if R₁ is O-R₃, then R₃ can be a fluorinated






5 moiety such as wherein p can be an integer from 0 to 3, and the F can be an ¹⁸F. In some configurations, p can be 2.


In various aspects of the above embodiments, a radiolabeled compound or salt thereof



can be a molecular species such as










The present teachings also include various analogues of dihydroethidium (DHE) that can serve as intermediates or precursors in syntheses of the radiolabeled compounds disclosed herein.

- 5 The present teachings also include methods of synthesis of radiolabeled compounds described herein, as well as synthesis of various intermediates and precursors.


The present teachings include, without limitation, the following aspects:

1. A radiolabeled compound or salt thereof of structure






, or , q is an integer from 0 to 10; n is an integer from 0 to 3, m is an integer from 0 to 3, and p is an integer from 0 to 3, R₂ is methyl or



, and a is an integer from 0 to 10, wherein at least one atom is a radioisotope.


- 5 2. A radiolabeled compound or salt thereof in accordance with aspect 1, wherein the radioisotope is a positron-emitting radioisotope.
- 3. A radiolabeled compound or salt thereof in accordance with aspect 1, wherein R₁ is H.
- 4. A radiolabeled compound or salt thereof in accordance with aspect 3, wherein the CH₃ is an ¹¹CH₃.
- 10 5. A radiolabeled compound or salt thereof in accordance with aspect 1, wherein R₃ is (CH₂)_q-CH₃ and q is 0.
- 6. A radiolabeled compound or salt thereof in accordance with aspect 5, wherein R₃ is ¹¹CH₃.
- 7. A radiolabeled compound or salt thereof in accordance with aspect 1, wherein R₃ is



, n is an integer from 0 to 3, and m is an integer from 0 to 3.

- 15 8. A radiolabeled compound or salt thereof in accordance with aspect 1, wherein R₁ is O-R₃, R₃ is CH₃.
- 9. A radiolabeled compound or salt thereof in accordance with aspect 8, wherein R₁ is O-R₃, wherein R₃ is ¹¹CH₃.

10. A radiolabeled compound or salt thereof in accordance with aspect 1, wherein R_1 is $O-R_3$,




n is an integer from 0 to 3 and m is an integer from 0 to 3, and F is ^{18}F .

11. A radiolabeled compound or salt thereof in accordance with aspect 10, wherein n is 1.

5 12. A radiolabeled compound or salt thereof in accordance with aspect 10, wherein m is 2.

13. A radiolabeled compound or salt thereof in accordance with aspect 10, wherein n is 1 and m is 2.


14. A radiolabeled compound or salt thereof in accordance with aspect 1, wherein R_3




p is an integer from 0 to 3 and F is ^{18}F .



10 15. A radiolabeled compound or salt thereof in accordance with aspect 13, wherein p is 2.


16. A radiolabeled compound or salt thereof in accordance with aspect 1, wherein the compound



is selected from the group consisting of














17. A method of imaging reactive oxygen species (ROS) distribution in a mammal, the method comprising:

- administering to the mammal a compound or salt thereof, of any one of aspects 1-15; and
- 5 subjecting the mammal to PET scanning.







NaBH₃CN.





21. A method of synthesizing , comprising contacting with EtOAc/HCl.




5 22. A method of synthesizing , comprising




contacting with 4-methoxybenzoyl chloride and chlorobenzene.







24. A method of synthesizing



, comprising






contacting

with HCOONH_4 , $\text{Pd}(\text{OH})_2/\text{C}$.












28. A method of synthesizing _____, comprising






29. A method of synthesizing _____, comprising




contacting _____ with HCOONH_4 and $\text{Pd}(\text{OH})_2/\text{C}$.



30. A method of synthesizing




_____, comprising contacting






31. A method of synthesizing











32. A method of synthesizing _____, comprising









































42. A method of synthesizing



, comprising














































55. A method of synthesizing _____, comprising









57. A method of synthesizing

, comprising contacting




with EtOAc/HCl

58. A method of synthesizing




comprising contacting




with (Boc)₂O.

59. A method of synthesizing







comprising contacting



with CH₃I,

5 followed by NaBH₃CN.



BRIEF DESCRIPTION OF THE DRAWINGS

5 FIG. 1 illustrates the fluorescence intensity over time in a platereader study of oxidation rate of compound **C1**. The X-axis scale is in hours:minutes:seconds. The fluorescence intensity is measured in arbitrary units (a.u.).

FIG. 2 presents the production of **C1** oxidation fluorescence signal compared to DHE for a variety of conditions.

10 FIG. 3 presents a comparison of compounds **WC-63**, **WC-77** and **WC-81** for selective reactivity with superoxide radical. In FIG. 3, the data for compounds **WC-63**, **WC-77** and **WC-81** are labeled PET63, PET77 and PET82, respectively.

FIG. 4 presents confocal microscope images of in vivo oxidation of compound **C1** in mouse brains pretreated with subanesthetic doses of ketamine and sacrificed 16 hours after administration of compound **C1**. In FIG. 4, panel G is an enlarged image of panel B.

15 FIG. 5 presents quantitative measures of fluorescence of oxidized forms of DHE and compound **C1** in mouse brain.

FIG. 6 presents quantitative measures of regional brain fluorescence of compound **WC-81** in the cortices and cerebellum of live mice.

20 FIG. 7 presents images of oxidation of compound **WC-81** in mouse brain. Panel A shows oxidation of compound **WC-81** localized in the extracellular space around cortical neurons. Panel B shows oxidation of compound **WC-81** localized in the extracellular space around hippocampal neurons. Panel C shows oxidation of compound **WC-81** localized in cerebral microvessels.

25 FIG. 8 presents images of fluorescence lifetimes for the oxidation products of compounds **WC-63**, **WC-77** and **WC-81** in the brain and chest. In FIG. 8, all images are calibrated to the same scale.

DETAILED DESCRIPTION


30 The present inventors disclose herein a series of compounds, including radiolabeled compounds, that can be used as tracers for imaging distribution of reactive oxygen species in a mammalian subject such as a human. Distribution of ROS using a compound disclosed herein as a tracer can be determined by fluorescence imaging, or by positron emission topography (PET) imaging. In some embodiments, a compound can comprise a radioisotope, such as a positron

emitter. Accordingly, a compound of the present teachings can comprise a radioisotope such as ^{18}F or ^{11}C .

In some aspects, the inventors provide methods of imaging ROS in tissue in a human or other animal subject such as a mammalian subject. These methods comprise administering to the subject a radiolabeled compound, and imaging distribution of the radiolabel by PET scanning. In some configurations, the PET scanning can yield an image which can then be interpreted by a medical professional, such as a physician.


Without limitation, a radiolabeled compound of the present teachings can be useful in imaging distribution of ROS in tissue of a human or mammal, such as, for example, imaging distribution of ROS in the brain of a human or mammal. Imaging distribution of ROS in the brain can be useful, for example, for better understanding the role of ROS in the pathogenesis of Alzheimer's disease, cerebral vascular disease, Parkinson's disease and schizophrenia and for developing therapies as well as monitoring response to therapy. Thus, the present radiotracers and methods can be used, for example, by a medical professional to determine if a therapy is effective.

In some embodiments, a compound of the present teachings can be a radiolabeled compound or salt thereof of structure



wherein R_1 is H or $\text{O}-R_3$, R_3 is $(\text{CH}_2)_q-\text{CH}_3$, $(\text{---})_n$, or $(\text{---})_p-\text{F}$, q is an integer from 0 to 10, n is an integer from 0 to 3, m is an integer from 0 to 3, and p is integer from 0 to 3.

In additional embodiments of the present teachings, the inventors disclose methods of imaging ROS distribution in a mammal such as a human. These methods comprise administering to the mammal a radiolabeled compound or salt thereof of structure



5 wherein R_1 is H or $O-R_3$, R_3 is $(CH_2)_q-CH_3$, $-(CH_2)_n-F$, or $-(CH_2)_p-F$, q is an integer from 0 to 10, n is an integer from 0 to 3, m is an integer from 0 to 3, and p is integer from 0 to 3, and subjecting the mammal to PET scanning.

In some embodiments, a compound of the present teachings can be a radiolabeled compound or salt thereof as disclosed herein, wherein R_1 is H.


10 In some embodiments, a compound of the present teachings can be a radiolabeled compound or salt thereof as disclosed herein, wherein R_1 is H and the CH_3 is $^{11}CH_3$.

In some embodiments, a compound of the present teachings can be a radiolabeled compound or salt thereof as disclosed herein, wherein R_1 is $O-R_3$, R_3 is $(CH_2)_q-CH_3$, q is 0.

15 In some embodiments, a compound of the present teachings can be a radiolabeled compound or salt thereof as disclosed herein, wherein R_1 is $O-R_3$, R_3 is CH_3 .


In some embodiments, a compound of the present teachings can be a radiolabeled compound or salt thereof as disclosed herein, wherein R_1 is $O-R_3$, R_3 is $^{11}CH_3$.

In some embodiments, a compound of the present teachings can be a radiolabeled compound or salt thereof as disclosed herein, wherein R_1 is $O-R_3$, wherein R_3 is




, wherein n is an integer from 0 to 3, m is an integer from 0 to 3, and F is ¹⁸F.

In some embodiments, a compound of the present teachings can be a radiolabeled compound or salt thereof as disclosed herein, wherein R₁ is O-R₃, wherein R₃ is




5 , wherein n is 1, m is an integer from 0 to 3, and F is ¹⁸F.

In some embodiments, a compound of the present teachings can be a radiolabeled compound or salt thereof as disclosed herein, wherein R₁ is O-R₃, wherein R₃ is





, wherein n is an integer from 0 to 3, m is 2, and F is ¹⁸F.

10 In some embodiments, a compound of the present teachings can be a radiolabeled compound or salt thereof as disclosed herein, wherein R₁ is O-R₃, wherein R₃ is





, wherein n is 1, m is 2 and F is ¹⁸F.



In some embodiments, a compound of the present teachings can be a radiolabeled compound or salt thereof as disclosed herein, wherein R_1 is $O-R_3$, wherein R_3 is , wherein p is an integer from 0 to 3 and F is ^{18}F .

In some embodiments, a compound of the present teachings can be a radiolabeled DHE analogue or salt thereof as disclosed herein, wherein R_1 is $O-R_3$, wherein R_3 is , wherein p is 2 and F is ^{18}F .


In various aspects of the above embodiments, a radiolabeled DHE analogue or salt thereof can include particular molecular species, such as




10




In further embodiments of the present teachings, the inventors disclose a radiolabeled compound or salt thereof of structure



wherein R_1 is H or $O-R_3$; R_3 is $(CH_2)_q-CH_3$,




, and wherein q is an integer from 0 to 10; n is an integer from 0 to 3; m is an integer from 0 to 3; p is an integer from 0 to 3; and a is an integer from 2 to 10. In some configurations, R_1 is H. In some configurations, CH_3 is $^{11}CH_3$. ϕ represents a phenyl group. In some configurations, R_3 is $(CH_2)_q-CH_3$ and $q=0$. In some configurations, R_3 is $^{11}CH_3$.




In some configurations, R_3 is , n is an integer from 0 to 3, and m is an integer from 0 to 3. In some configurations, R_1 is $O-R_3$, and R_3 is $^{11}CH_3$.

In some configurations, the radiolabeled compound or salt thereof is of structure




, wherein a is an integer from 2 to 10. In some

configurations, the radiolabeled compound or salt thereof is of structure




, wherein a is an integer from 2 to 10. In some

configurations, the radiolabeled compound or salt thereof is of structure




, wherein a is an integer from 2 to 10. In some

5 configurations, the radiolabeled compound or salt thereof is of structure



, wherein Z is (CH₂)_q-CH₃, q is an integer




from 0 to 10,

, or

, and a is an integer from 2 to 10.


In some configurations, the radiolabeled compound or salt thereof is of



structure

, wherein a is an integer from 2 to 10


and q is an integer from 0 to 10. In some configurations, the CH₃ can be ¹¹CH₃. In some configurations, the radiolabeled compound or salt thereof is of structure



5


, wherein a is an integer from 2 to 10. In some

configurations, the radiolabeled compound or salt thereof is of structure




, wherein a is an integer from 2 to 10. In some


configurations, the radiolabeled compound or salt thereof is of structure




n is an integer from 0 to 3, and m is an integer from 0 to 3. In some configurations, the radiolabeled compound or salt thereof is of structure



5 10, n is an integer from 0 to 3, and m is an integer from 0 to 3. In some configurations, the radiolabeled compound or salt thereof is of structure



integer from 0 to 3. In some configurations the selected compound or salt thereof is of



structure

, wherein a is an integer from 2 to 10

and p is an integer from 0 to 3.

In various aspects of the embodiments, methods for synthesis of the compounds disclosed herein are provided. In particular aspects, methods of synthesis of radiolabeled compounds or salts thereof are provided. In further aspects, methods for synthesis of radiolabeled DHE analogue precursors are also provided.


EXAMPLES

The following examples are illustrative of the various embodiments of the present teachings. The following examples provide non-limiting illustrations of the present teachings. While some of examples may include conclusions about the way the invention may function, the inventor do not intend to be bound by those conclusions, but put them forth only as possible explanations. Unless noted by use of past tense, presentation of an example does not imply that an experiment or procedure was, or was not, conducted, or that results were, or were not actually obtained. The examples are not intended to limit the scope of any claim.

The methods described herein utilize laboratory techniques well known to skilled artisans, and guidance can be found in laboratory manuals and textbooks such as Sambrook, J., et al., *Molecular Cloning: A Laboratory Manual*, 3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 2001; Spector, D. L. et al., *Cells: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1998; and Harlow, E., *Using Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1999; Hedrickson et al., *Organic Chemistry* 3rd edition, McGraw Hill, New York, 1970; Carruthers, W., and Coldham, L., *Modern Methods of Organic Synthesis* (4th Edition), Cambridge University Press, Cambridge, U.K., 2004; Curati, W.L., *Imaging in Oncology*, Cambridge University Press, Cambridge, U.K., 1998; Welch, M.J., and Redvanly, C.S., eds. *Handbook of Radiopharmaceuticals: Radiochemistry and Applications*, J. Wiley, New York, 2003.


Example 1

This example illustrates synthesis of compound **C1**, as shown below.




5 Example 2


This example illustrates synthesis of compound **WC-63**, as shown below.



Example 3


This example illustrates synthesis of compound WC-77, as shown below.





Example 4

This example illustrates synthesis of compound **WC-81**, as shown below.




5

Example 5


This example illustrates radiosynthesis of compound [¹¹CH₃]Cl, as shown below.


Scheme 1:



5

Scheme 2:






Two different routes for making [¹¹C]C1 are used. The first route (Scheme 1) involves N-alkylation of the phenanthridine analog with [¹¹C]methyltriflate followed by reduction of the nitrogen-carbon double bond with sodium borohydride. Deprotection with HCl results in the formation of [¹¹C]C1. The second approach (Scheme 2) begins with the dihydrophenanthridine analog is then be labeled with [¹¹C]methyltriflate and removal of the butyloxycarbonyl (Boc) protecting groups results in the formation of [¹¹C]C1.

Example 6


This example illustrates a synthesis of radiolabeled compound [¹¹C]WC-63, as shown below.




In this example, synthesis of compound [¹¹C]WC-63 involves reaction of a phenol precursor with [¹¹C]methyl iodide in the presence K₂CO₃, followed by removal of the Boc protecting groups with EtOAc/HCl.

15 Example 7


This example illustrates radiosynthesis of compound [¹⁸F]WC-77, as shown below.





Example 8

This example illustrates radiosynthesis of compound [¹⁸F]WC-81, as shown below.



5

Example 9

This example illustrates oxidation rates of Compound **C1** under various conditions. In these experiments, fluorescence intensity of Compound **C1** was evaluated in a plate reader assay, and the fluorescence intensity over time of the oxidation rate of compound **C1** was evaluated. As illustrated in FIG. 1, compound **C1** showed both reactivity and selectivity for superoxide. Excitation was done at 480 nm and emission filtering was > 590 nm. The following reaction conditions were used:

10

CI	158 μ M
XO	0.052 U/ml
hypoxanthine	1mM
SOD	575 U/ml
Catalase	60 U/ml
Hydrogen peroxide	1 mM
HRP	0.2 U/ml

Under the conditions of superoxide production with hypoxanthine and xanthine oxidase (XO system), compound **CI** is oxidized rapidly. Compound **CI** showed no oxidation with peroxide (H_2O_2), or H_2O_2 with horseradish peroxidase. Superoxide dismutase (SOD), which rapidly removes superoxide, also completely removed compound **CI** oxidation. Finally, addition of catalase, which removes H_2O_2 , did not effect compound **CI** oxidation, demonstrating independence from H_2O_2 . The rate of conversion by oxidation of **CI** was slightly faster than DHE itself (data not shown) and was selective for superoxide as there was no observable oxidation from H_2O_2 or H_2O_2 in the presence of horseradish peroxidase (to generate the hydroxyl radical, HO \cdot).

Example 10

This example illustrates comparative oxidation of **CI** and DHE. In these experiments, fluorescence of **CI** or DHE were determined in the presence of oxidizers. As shown in FIG. 2, **CI** showed selective oxidation similar to DHE with elevations in the presence of superoxide but not in the presence of peroxide (H_2O_2), hydroxyl radical (HO \cdot) or peroxynitrate (ONOO \cdot). In this single wavelength fluorescence plater reader study, it should be noted that the ratio of fluorescence increase in a study cannot be compared between DHE and compound **CI** since their non-oxidized forms have fluorescence and the light spectra overlap to a variable degree with the oxidized forms.

20 Example 11

This example illustrates a comparison of compounds for selective reactivity with superoxide radical. In these experiments, compounds **WC-63**, **WC-77** and **WC-81** were exposed to superoxide generated by xanthine oxidase/hypoxanthine in the absence or presence SOD, and were evaluated similarly as compound **C1** in Example 10. The concentration of DHE for all experiments was double that of compounds **WC-63**, **WC-77** and **WC-81**. As shown in FIG. 3, compounds **WC-63**, **WC-77** and **WC-81** showed high specificity for superoxide with generally no evidence of oxidation with H₂O₂ or hydroxyl radical. The exception was compound **WC-81**, which had partial oxidation in the presence of the hydroxyl radical at these supraphysiologic levels, but remained intact. Change in fluorescent units from baseline is shown on the y-axis under same conditions as FIG. 2. The data showed that DHE, **WC-63**, **WC-77** and **WC-81** were each oxidized by superoxide, and that SOD fully blocked oxidation, indicating that oxidation is superoxide-dependent. All of the compounds were also exposed to H₂O₂ in the presence of horseradish peroxidase (HRP) to generate hydroxyl radical. H₂O₂ alone did not oxidize any of the compounds. A second method of generating superoxide through SIN-1 decomposition (Sydnominine-1, an extracellular donor of NO and superoxide, Panagiotidis, G., et al., Br. J. Pharmacol. 114: 289-296, 1995) also indicated that all 4 compounds detect superoxide, and SOD was able to block oxidation completely. Compound **WC-77** was the most specific compound for superoxide, but compound **WC-81** was the most stable and intact product. It was still highly selective for superoxide under our experimental conditions, in which the levels of hydroxyl radical were very supraphysiological.

20 Example 12

This example illustrates imaging of oxidation of Compound **C1** in an animal model.

In these experiments, mice were pretreated for several days with subanesthetic doses of ketamine, which produces a large increase in neuronal superoxide generation (Behrens et al, 2007). The mice were injected with 27 mg/kg of compound **C1** and sacrificed 16 hours later. Brains were sliced and imaged by confocal microscopy. The images of the brains sliced 16 hours after administration of compound **C1** are depicted in FIG. 4. The images show a pattern of fluorescence consistent with the presence of intracellular oxidized compound **C1**. As shown in FIG. 5, both DHE and compound **C1** detected a large increase in superoxide produced by ketamine treatment. The quantitative measures of fluorescence of the oxidized forms of DHE and compound **C1** in mouse brain are depicted in FIG. 5. When compound **C1** was administered to transgenic mice overexpressing SOD, the increased fluorescence from the ketamine pretreatment was abolished, consistent with the SOD removal of superoxide (FIG. 5).

Example 13

This example illustrates kinetics of Compound **WC-81** in brain in vivo.

In these experiments, an anesthetized mouse was placed in an eXplore Optix™-MX2 scanner, and an initial scan was completed for background fluorescence (T=0). A single
5 intraperitoneal injection of 50mg/kg of compound **WC-81** in 50% DMSO, 50% saline was performed, and the animal was rescanned at 10 minutes. It took approximately 5 minutes to complete the scan. The mouse was removed, allowed to recover, then re-anesthetized and rescanned at 1 hour. A similar recovery and rescan was performed at 2 hours. The mouse was pretreated with subanesthetic doses of ketamine to raise superoxide levels. Cortical (left and
10 right) and cerebellar activity are depicted in FIG. 6. The results show that compound **WC-81** penetrated the brain. While a plateau was reached by 60 minutes, there was a fall in levels at 120 minutes.

Example 14

This example illustrates fluorescence from oxidation of Compound **WC-81**. In these
15 experiments, compound **WC-81** was administered to mice, and images of mouse brains were then obtained (FIG. 7). Fluorescence from oxidation of compound **WC-81** was localized to cerebral microvessels and to the extracellular space around certain cortical or hippocampal neurons.

Example 15



20 This example illustrates tissue uptake of the tracers. In these experiments, tracer lifetimes in mice were studied by fluorescence imaging of DHE and compounds **WC-63**, **WC-77** and **WC-81**. As shown in FIG. 8, DHE, compounds **WC-63** and **WC-81** are present in the brain and organs in the chest.

All references cited herein are incorporated by reference, each in its entirety.


CLAIMS

What is claimed is:

1. A radiolabeled compound or salt thereof of structure



5 wherein R_1 is H or $O-R_3$, R_3 is $(CH_2)_qCH_3$, or $(CH_2)_p$, q is an integer from 0 to 10; n is an integer from 0 to 3, m is an integer from 0 to 3, and p is an integer



from 0 to 3, R_2 is methyl or $(CH_2)_n$, and a is an integer from 0 to 10, wherein at least one atom is a radioisotope.


2. A radiolabeled compound or salt thereof in accordance with claim 1, wherein the radioisotope is a positron-emitting radioisotope.

3. A radiolabeled compound or salt thereof in accordance with claim 1 or claim 2, wherein R_1 is H.

4. A radiolabeled compound or salt thereof in accordance with any one of claims 1-3, wherein the compound or salt thereof comprises a methyl, and wherein the methyl is an $^{11}CH_3$.

5. A radiolabeled compound or salt thereof in accordance with claim 1, wherein R_3 is $(CH_2)_qCH_3$ and q is 0.

6. A radiolabeled compound or salt thereof in accordance with claim 5, wherein R_3 is $^{11}\text{CH}_3$.
7. A radiolabeled compound or salt thereof in accordance with claim 1 or claim 2, wherein R_3 is




, n is an integer from 0 to 3, and m is an integer from 0 to 3.

8. A radiolabeled compound or salt thereof in accordance with claim 1 or claim 2, wherein R_1 is $\text{O}-R_3$, R_3 is CH_3 .

9. A radiolabeled compound or salt thereof in accordance with claim 8, wherein R_1 is $\text{O}-R_3$, wherein R_3 is $^{11}\text{CH}_3$.

10. A radiolabeled compound or salt thereof in accordance with claim 1 or claim 2, wherein R_1




is $\text{O}-R_3$, R_3 is , n is an integer from 0 to 3, m is an integer from 0 to 3, and F is

10 ^{18}F .




11. A radiolabeled compound or salt thereof in accordance with claim 10, wherein n is 1.
12. A radiolabeled compound or salt thereof in accordance with claim 10, wherein m is 2.
13. A radiolabeled compound or salt thereof in accordance with claim 10, wherein n is 1 and m is 2.

- 15 14. A radiolabeled compound or salt thereof in accordance with claim 1 or claim 2, wherein R_3



is , p is an integer from 0 to 3 and F is ^{18}F .

15. A radiolabeled compound or salt thereof in accordance with claim 13, wherein p is 2.
16. A radiolabeled compound or salt thereof in accordance with claim 1 or claim 2, wherein the compound is selected from the group consisting of








FIG. 1

- — X0 SYSTEM MINUS HYPOXANTHINE
- X0 SYSTEM
- ▼ --- X0 SYSTEM PLUS CATALASE
- ▽ --- X0 SYSTEM PLUS SOD
- --- HYDROGEN PEROXIDE
- --- HYDROGEN PEROXIDE PLUS HRP




2/11

FIG. 2



3/11

FIG. 3



4/11

FIG. 4A

FIG. 4B

FIG. 4C




FIG. 4D

FIG. 4E

FIG. 4F






FIG. 4G


5/11

FIG. 5



6/11

FIG. 6



7/11

FIG. 7A




FIG. 7B





FIG. 7C




8/11


FIG. 8

COLOR IMAGE

BRAIN



CHEST




9/11


FIG. 8

BLUE CHANNEL

BRAIN



CHEST




10/11


FIG. 8

GREEN CHANNEL

BRAIN



CHEST




11/11

FIG. 8

RED CHANNEL

BRAIN



CHEST

